



Landtype Association Descriptions for the Superior Coastal Plain Ecological Landscape		
212lc01	Apostle Islands	The characteristic landform pattern is variable modified lacustrine moraine and lake shore. Soils are predominantly well drained clay loam, and sand over clay or loam till, outwash, or sandstone. Common habitat types include ASnMi, AbArSn, hydromesic, ATM, PArVAa-Po and forested lowland.
212Ya01	Douglas Lake-Modified Till Plain	The characteristic landform pattern is undulating modified lacustrine moraine with deep v-shaped ravines. Soils are predominantly somewhat poorly drained clay over calcareous clay till or loamy lacustrine. Common habitat types include AbArSn, forested lowland and ASnMi.
212Ya02	Bayfield Lake-Modified Till Plain	The characteristic landform pattern is hilly modified lacustrine moraine with deep v-shaped ravines. Soils are predominantly well drained complex of clay, loam, and sand over clayey or loamy till, or outwash. Common habitat types include ASnMi, AbArSn, hydromesic, ATM, PArVAa-Po and forested lowland.
212Ya03	Ashland Lake-Modified Till Plain	The characteristic landform pattern is undulating modified lacustrine moraine with deep v-shaped ravines. Soils are predominantly somewhat poorly drained clay loam over calcareous clay till or loamy lacustrine. Common habitat types include AbArSn, ASnMi, hydromesic and forested lowland.
212Ya04	Iron Gogebic Dissected Lake Bench	The characteristic landform pattern is undulating modified lacustrine moraine with deep v-shaped ravines. Soils are predominantly somewhat poorly drained silt loam over calcareous silty clay loam till or loamy lacustrine. Common habitat types include AbArSn, hydromesic, ASnMi and AOCa.
212Ya07	Bibon Marsh	The characteristic landform pattern is nearly level swamp with outwash plains and alluvial plains common. Soils are predominantly very poorly drained organic deposits. Common habitat type is forested lowland.
212Ya08	Kakagon/Bad River Sloughs	The characteristic landform pattern is nearly level delta and outwash plains. Soils are predominantly very poorly drained organic deposits and sandy to clayey alluvium. Common habitat type is forested lowland.
212Ya09	Carlton Plains	The characteristic landform pattern is undulating outwash plains and lake plains with old beaches and dunes common. Soils are predominantly excessively drained sand over outwash or loamy lacustrine. Common habitat types include hydromesic, ArAbVCo, AVCI, ASnMi, PArV-U and forested lowland.