

**LTA: 212Ob03**NAME:  
Chambers IslandsACRES:  
3003.208SQUARE MILES:  
4.693**DESCRIPTION:**

The characteristic landform pattern is gently sloping with beaches common. Soils are predominantly sandy. Common habitat types include AFSt, FArAa, forested lowland, and TMC.

**CLIMATE**CODE                      PERCENT  
79                              85**GEOLOGY**BEDROCK TYPE DESCRIPTION  
SandstoneAVERAGE DEPTH  
TO BEDROCK  
10000BEDROCK DEPTH  
DESCRIPTION  
Bedrock is between 100 feet and 50 feet of the land surface**GEOMORPHOLOGY**GEOMORPHOLOGY PROCESS  
Lake DepositionTOPOGRAPHY  
UndulatingSURFACE  
Beach**SOIL INFORMATION**SOIL ASSOCIATIONS  
Rousseau-Kiva-Markey**SOIL DESCRIPTION**

Well drained and moderately well drained sandy soils with a fine sand or fine sandy loam surface over non-calcareous or calcareous fine sand beach deposits, along with very poorly drained nonacid organic soils.

SURFACE TEXTURES  
FS-FSL-MKGENERAL TEXTURES  
Sandy-MuckyFAMILY TEXTURES  
S-SPDRAINAGE CLASSES  
WD-MWD-VPDPARENT MATERIAL  
Beach-Organic**KOTAR'S HABITAT**

HABITAT 1	HABITAT 2	HABITAT 3	HABITAT 4	HABITAT 5	HABITAT 6
AFSt	FArAa	Lowland	TMC		

\*Listed in order of probability occurrence, with each having an occurrence of 10% or greater

**WISCLAND LAND COVER**

COVER TYPE CLASS	ACRES	PERCENT
Agricultural Land	285	9
Bare Land	9	0
Forested Wetland	52	2
Grassland	64	2
Nonforested Wetland	23	1
Open Water	375	13
Upland Broad-leaved Deciduous	1964	66
Upland Coniferous Forest	224	7

**FINLEY'S PRESETTLEMENT VEGETATION INFORMATION**

CODE	PERCENT	DEFINITION
Hydrographic	12	Water
BHH/P	63	Beech, Hemlock, Sugar Maple, Yellow Birch, White Pine, Red Pine
HH/P	0	Hemlock, Sugar Maple, Yellow Birch, White Pine, Red Pine
PW/PR	14	White Pine, Red Pine

**HYDROLOGY**

PERENNIAL STREAMS	INTERMITTENT STREAMS	OPEN WATER	MARSH ACRES	DEPTH TO AQUIFER	SURFACE DRAINAGE
0 Miles	0 Miles	352 Acres	2991 Acres	0'-20'	dendritic